

Citation for published version:

Gonzalez, JT, Fuchs, CJ, Smith, FE, Thelwall, PE, Taylor, R, Stevenson, EJ, Trenell, MI, Cermak, NM & van Loon, LJC 2015, 'Ingestion of glucose or sucrose prevents liver but not muscle glycogen depletion during prolonged endurance-type exercise in trained cyclists', *American Journal of Physiology: Endocrinology and Metabolism*, vol. 309, no. 12, pp. E1032-E1039. <https://doi.org/10.1152/ajpendo.00376.2015>

DOI:

[10.1152/ajpendo.00376.2015](https://doi.org/10.1152/ajpendo.00376.2015)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

©American Psychological Association, 2015. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1152/ajpendo.00376.2015](https://doi.org/10.1152/ajpendo.00376.2015)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Ingestion of Glucose or Sucrose Prevents Liver but not Muscle
Glycogen Depletion During Prolonged Endurance-type Exercise in
Trained Cyclists**

Javier T. Gonzalez^{1,2}, Cas J. Fuchs^{2,3}, Fiona E. Smith⁴, Pete E. Thelwall⁴, Roy
Taylor⁴, Emma J. Stevenson², Michael I. Trenell⁴, Naomi M. Cermak³, and
Luc J.C. van Loon³

*¹Department for Health, University of Bath, Bath, United Kingdom; ²Faculty of
Health and Life Sciences, Northumbria University, Newcastle-upon-Tyne,
United Kingdom; ³Department of Human Movement Sciences, NUTRIM
School of Nutrition and Translational Research in Metabolism, Maastricht
University Medical Centre+ (MUMC+), Maastricht, The Netherlands;
⁴Newcastle Magnetic Resonance Centre, Institute of Cellular Medicine,
Newcastle University, Newcastle-upon-Tyne, United Kingdom.*

Running head: Carbohydrate ingestion and liver glycogen depletion

Corresponding author

Luc J.C. van Loon

Department of Human Movement Sciences, NUTRIM School of Nutrition and
Translational Research in Metabolism, Maastricht University Medical Centre+
(MUMC+), Maastricht, The Netherlands.

E-mail: l.vanloon@maastrichtuniversity.nl

ABSTRACT

Purpose: To define the effect of glucose ingestion compared to sucrose ingestion on liver and muscle glycogen depletion during prolonged endurance-type exercise. **Methods:** Fourteen cyclists completed two 3-h bouts of cycling at 50% of peak power output while ingesting either glucose or sucrose at a rate of 1.7 g/min (102 g/h). Four cyclists performed an additional third test in which only water was consumed for reference. We employed ^{13}C magnetic resonance spectroscopy to determine liver and muscle glycogen concentrations before and after exercise. Expired breath was sampled during exercise to estimate whole-body substrate use. **Results:** Following glucose and sucrose ingestion, liver glycogen levels did not show a significant decline following exercise (from 325 ± 168 to 345 ± 205 and 321 ± 177 to 348 ± 170 mmol/L, respectively; $P > 0.05$) with no differences between treatments. Muscle glycogen concentrations declined (from 101 ± 49 to 60 ± 34 and 114 ± 48 to 67 ± 34 mmol/L, respectively; $P < 0.05$), with no differences between treatments. Whole-body carbohydrate utilization was greater with sucrose (2.03 ± 0.43 g/min) vs glucose ingestion (1.66 ± 0.36 g/min; $P < 0.05$). Both liver (from 454 ± 33 to 283 ± 82 mmol/L; $P < 0.05$) and muscle (from 111 ± 46 to 67 ± 31 mmol/L; $P < 0.01$) glycogen concentrations declined during exercise when only water was ingested. **Conclusion:** Both glucose and sucrose ingestion prevent liver glycogen depletion during prolonged endurance-type exercise. Sucrose ingestion does not preserve liver glycogen concentrations more than glucose ingestion. However, sucrose ingestion does increase whole-body carbohydrate utilization compared to glucose ingestion. This trial was

50 registered at clinicaltrials.gov as NCT02110836. **Keywords:** glucose; hepatic;
51 metabolism; nutrition; sucrose

Introduction

Carbohydrate and fat are the main substrates oxidized during moderate-intensity, endurance-type exercise (41). In the fasted state, muscle glycogen and plasma glucose are predominant sources of carbohydrate for oxidation (41), the latter continuously replenished by glycogenolysis and gluconeogenesis from the liver, with smaller contributions from the kidneys and intestine (30). Consequently, in the absence of carbohydrate consumption, liver and muscle glycogen concentrations decrease by 40-60% within 90 min of exercise at a workload of 70% of peak oxygen uptake ($\dot{V}O_{2\text{ peak}}$) (6, 37). Given the importance of liver glycogen for metabolic regulation (16), and the close relationship between liver glycogen content and exercise tolerance (6), it is important to understand the impact of carbohydrate ingestion on liver glycogen depletion during exercise.

Carbohydrate feeding during prolonged (>2 h) moderate-to-high intensity, endurance-type exercise enhances endurance performance and capacity (42), attributed to the facilitation of high rates of carbohydrate oxidation, prevention of hypoglycaemia and (under certain conditions) sparing of muscle glycogen (7, 38). Though some support has been provided that carbohydrate ingestion can attenuate muscle glycogen depletion (36, 39, 40), others have failed to confirm these findings (8, 12, 15, 20). Furthermore, prevention of liver glycogen depletion has been suggested (3, 20, 21), but this has never been experimentally assessed. We speculate that carbohydrate ingestion during exercise attenuates the decline in both liver as well as skeletal muscle glycogen contents.

To maximize carbohydrate availability during exercise, carbohydrate digestion and absorption should be optimized. Previous work suggests that exogenous glucose uptake by the gastrointestinal tract during exercise is restricted to ~1 g/min (5, 17, 19), attributed to saturation of the sodium-glucose luminal transporter-1 (SGLT-1). However, combined ingestion of glucose and fructose at ≥ 1.8 g/min has been shown to result in much higher exogenous carbohydrate oxidation rates (up to 1.75 g/min), compared to the ingestion of equal amount of glucose alone (17, 19). The greater uptake and oxidative capacity of glucose and fructose mixtures has been attributed to fructose being absorbed by the glucose transporter-5 (GLUT-5) in the intestine (11). As sucrose (commonly referred to as table sugar) combines glucose and fructose monomers, and sucrose hydrolysis is not rate limiting for intestinal absorption (14, 43), we hypothesize that sucrose ingestion at a rate exceeding 1 g/min will enhance exogenous carbohydrate availability when compared to the ingestion of an isoenergetic amount of glucose or glucose polymers. Moreover, since fructose appears to be preferentially directed to liver glycogen storage (relative to glucose) (32), sucrose may further prevent liver glycogen depletion during exercise.

The present study aimed to investigate the effect of high rates of glucose and sucrose ingestion on net changes in liver and muscle glycogen contents and intramyocellular lipid concentrations using magnetic resonance spectroscopy (MRS). We hypothesized that high-rates of carbohydrate ingestion would spare liver glycogen during prolonged exercise, and that sucrose ingestion would better maintain liver glycogen relative to glucose ingestion.

Methods

Study design

Participants completed preliminary testing prior to 2 main trials, during which subjects either ingested glucose (GLU) or sucrose (SUC) in a randomized, double-blind, crossover design separated by 7-14 d. Trials were conducted at the Newcastle Magnetic Resonance Centre (Newcastle-upon-Tyne, UK) in accordance with the Second Declaration of Helsinki, and following approval from the Northumbria University Faculty of Health and Life Sciences Ethics Committee. Randomization was performed using online statistical software (<http://www.randomizer.org/>). Blinding and preparation of the test-drinks was performed by an assistant who was not involved in the exercise tests.

In addition to the two main trials, four participants completed an additional control trial (CON) as a reference to establish the change in liver glycogen concentration without carbohydrate ingestion. This was identical to the SUC and GLU trials, with the exception that only water was ingested during exercise (identical volume to GLU and SUC trials), and blood sampling was not performed.

Participants

Fifteen trained cyclists were recruited for the study. Inclusion criteria included healthy, endurance trained, male cyclists; $\dot{V}O_{2peak} \geq 50$ mL/min/kg/. Exclusion criteria included the use of medication that could influence substrate metabolism, smokers and any known metabolic disorders. One participant was unable to complete the full 3 h cycling protocol due to nausea on the GLU

trial and was therefore excluded from the analysis. Consequently, 14 participants completed the two main trials.

Preliminary testing

An incremental cycling test was performed on an electromagnetically braked cycle ergometer (Velotron, RacerMate Inc., Seattle, WA, USA) to determine peak power output (W_{peak}) and peak oxygen uptake ($\dot{V}O_{2\text{peak}}$). Following a 5-min warm-up at 100 W, the workload began at 150 W and was increased by 50 W every 2.5 min to voluntary exhaustion (23). Expired gas was sampled continuously to determine oxygen uptake (Oxycon gas analyser, CareFusion corporation, San Diego, CA, USA).

Main trials

Participants arrived at Newcastle Magnetic Resonance Centre at 0700-0800 h following a 12 h fast. Strenuous exercise was prohibited for 24 h prior to trials, and participants were asked to record and replicate dietary intake for 24 h prior to trials. The final meal consumed on the evening before the main trials was provided by the investigators to participants to standardize the macronutrient intake across participants for this meal (25 g protein, 51 g carbohydrate and 32 g fat; 2479 kJ; 592 kcal).

MRS was used to determine liver and muscle glycogen and intramyocellular lipid concentrations prior to and following 3 h of cycling. Following a 5 min warm-up at 100 W, power output was increased to 50% W_{peak} for the remainder of the 3 h. Immediately prior to exercise, participants were provided

with 600 mL (86.4 g carbohydrate) of the test-drink, and then a further 150 mL (21.6 g carbohydrate) every 15 min during exercise. Four of the 14 cyclists did not manage to consume all of the carbohydrate on their first trial (which was a GLU trial for two participants, and a SUC trial for the other two participants) and therefore their carbohydrate intake was replicated for their second trial (the carbohydrate intakes for these four cyclists therefore ranged from 238-281 g, mean \pm SD: 292 ± 101 g, compared to the 324 g prescribed). This led to an average rate of carbohydrate intake for the entire group of 1.7 ± 0.2 g/min and 1.7 ± 0.2 g/min (102 ± 12 g/h and 102 ± 12 g/h) during GLU and SUC trials, respectively ($P > 0.05$; **Table 1**).

Carbohydrate drinks

Carbohydrate drinks were prepared by mixing 108 g of carbohydrate with 750 mL of water in an opaque bottle. This was replicated two more times to produce 3 bottles, each with 750 mL of a 7% carbohydrate solution; 324 g of carbohydrate in total. Both sources of carbohydrate were from plants that use C_3 carbon fixation to minimize differences in the natural abundance of ^{13}C (33). Accordingly, the glucose drink was produced with dextrose monohydrate obtained from wheat (Roquette, France) and the sucrose drink was produced with granulated sugar beet (AB Sugar, UK).

Blood sampling and analysis

Prior to exercise, an intravenous catheter was inserted into an antecubital vein for regular sampling. Blood samples were obtained prior to the exercise bout, and every 30 min during exercise. Briefly, 10 mL of blood was collected

in EDTA-vacutainers and immediately centrifuged at 2000 g for 10 min at 4°C. Plasma was then aliquoted and stored at -80°C for subsequent determination of insulin (IBL International, Hamburg, Germany) and non-esterified fatty acid (NEFA) concentrations (WAKO Diagnostics, Richmond, VA) in duplicate (intra- and inter-assay coefficients of variation all <10%). An additional 20 µL of whole blood was collected in a capillary tube and was used to determine glucose and lactate concentrations immediately (Biosen C_line, EKF Diagnostics, Magdeberg, Germany).

Expired gas analysis

Expired breath samples were taken every 30 min throughout exercise using the Douglas bag technique (9) accounting for variance in ambient oxygen and carbon dioxide concentrations (1). A mouthpiece connected to a two-way, non-rebreathing valve (model 2730, Hans Rudolph, Kansas City, Missouri), was used to collect gas samples (60 s sample after a 60 s stabilization phase), analysed for concentrations of oxygen and carbon dioxide using a paramagnetic and infrared transducers, respectively (Servomex 5200S, Crowborough, East Sussex, UK). Sensors were turned on 60 min prior to a two-point calibration (zero: 100% nitrogen; span: 20% oxygen and 8% carbon dioxide) using accuracy certified gas standards (BOC Industrial Gases, Linde AG, Munich, Germany).

Ambient temperature, humidity and barometric pressure using a Fortin barometer (F.D. and company, Watford, UK) were recorded, and expired gas samples were corrected to standard temperature and pressure (dry). Volume

and temperature of expired gas samples were determined using a dry gas meter (Harvard Apparatus, Edenbridge, Kent, UK) and thermistor (model 810-080, ETI, Worthing, UK), respectively, during gas evacuation. Calibration of the dry gas meter was performed regularly with a 3-L syringe (Series 5530, Hans-Rudolph Inc, Kansas City, Missouri, USA).

Subjective ratings

Ratings of gut discomfort were assessed every 30 min during exercise using a 5-point scale, where 1 was anchored at “no discomfort” and 5 at “maximum discomfort”. Ratings of perceived exertion (RPE) were assessed using the Borg scale (2).

Measurement of muscle and liver glycogen

Tissue glycogen concentration was determined from the magnitude of the natural abundance signal from the C-1 carbon of glycogen at a frequency of 100.3 ppm. A Philips 3 Tesla Achieva scanner (Philips Healthcare, Best, The Netherlands) was used with a 6 cm diameter ^{13}C surface coil with integral ^1H decoupling surface coil (PulseTeq, Worton under Edge, UK) to measure muscle glycogen concentration and an in-house built 12 cm $^{13}\text{C}/^1\text{H}$ surface coil used to measure liver glycogen concentration.

For muscle glycogen measurements, the surface coil was placed over the widest part of the *Vastus lateralis* and the leg was held in position with fabric straps to prevent movement. Pulse power was calibrated to a nominal value of 80° by observing the power dependent variation in signal from a fiducial

marker located in the coil housing, containing a sample exhibiting ^{13}C signal with short T_1 (213 mM $[2\text{-}^{13}\text{C}]\text{-acetone}$ and 25 mM GdCl_3 in water). Automated shimming was carried out to ensure that the magnetic field within the scanner was uniform over the active volume of the ^{13}C coil. The ^{13}C spectra were acquired over 15 min using a non-localized ^1H decoupled ^{13}C pulse-acquire sequence (TR 120 ms, spectral width 8 kHz, 7000 averages, WALTZ decoupling). ^1H decoupling was applied for 60% of the ^{13}C signal acquisition to allow a relatively fast TR of 120 ms to be used within the Specific Absorption Rate safety limitations.

For liver glycogen measurements the $^{13}\text{C}/^1\text{H}$ surface coil was placed over the right lobe of the liver. Spectra were acquired over 15 min using non-localized ^1H decoupled ^{13}C pulse acquisition sequences (TR 300 ms, spectral width 8 kHz, 2504 averages, WALTZ decoupling, nominal ^{13}C tip angle of 80°). Scout images were obtained at the start of each study to confirm optimal coil position relative to the liver.

Tissue glycogen concentrations were calculated from the amplitude of the C1-glycogen ^{13}C signal using Java Based Magnetic Resonance User Interface (jMRUI) version 3.0 and the AMARES algorithm [7] as described in detail previously (25, 26, 28, 37).

Measurement of intramyocellular lipid

Intramyocellular lipid content was determined routinely, as described in more detail previously (37). In short, a 12 cm ^1H transmitter/receiver coil was used

to obtain ^1H spectra to measure intramyocellular lipid (IMCL) content in the widest part of the *gastrocnemius*. The PRESS (Point Resolved Spectroscopy) (4) sequence was used to acquire ^1H spectra from a $2 \times 2 \times 2$ cm voxel, using an echo time of 25 ms, spectral resolution of 1 Hz and repetition time of 5000 ms with 32 acquisitions. Spectra were analyzed with JMRUI version 3.0 using the least square fitting AMARES algorithm (4, 31). The inter-observer bias was 0.09 mmol/L with a 95% limit of agreement of 0.8 mmol/L ($P > 0.05$).

Calculations and statistical analysis

Due to the lack of data regarding exercise-induced changes in liver glycogen concentrations with carbohydrate feeding, a sample size estimation was based on data from exogenous carbohydrate oxidation rates (as a surrogate for carbohydrate availability). Sucrose increases exogenous carbohydrate oxidation during cycling by ~30% (18). Using this figure, along with the 7% intra-individual coefficient of variation of hepatic glycogen content measured by ^{13}C MRS (34), the study was designed to provide statistical power above 90% with an alpha level of 0.05 with a minimum sample size of $n = 7$ in a crossover design (10).

Whole-body rates of carbohydrate and lipid utilization were estimated using the following equations assuming negligible protein oxidation (13, 22):

$$\text{Net lipid utilization (g/min)} = (1.695 \times \dot{V}\text{O}_2) - (1.701 \times \dot{V}\text{CO}_2)$$

275 Net carbohydrate utilization (g/min) = $(4.210 \times \dot{V}CO_2) - (2.962 \times \dot{V}O_2)$

277 Units of $\dot{V}O_2$ and $\dot{V}CO_2$ are L/min

279 Unless otherwise stated, all data were expressed in the text as the mean \pm
280 standard deviation (SD) of the mean and the error bars presented in figures
281 are 95% confidence intervals (CI). Data were checked for normal distribution
282 and log-transformed if appropriate prior to statistical analysis.

284 Liver and muscle glycogen, and IMCL concentrations from the four
285 participants who completed the CON trial were assessed by two-way (trial x
286 time) repeated measures ANOVA with trial (GLU vs SUC vs CON) and time
287 (pre- vs post-exercise) as within-subject factors. Rates of substrate utilization
288 were assessed by a one-way repeated measures (GLU vs SUC vs CON)
289 ANOVA. No further inferential statistics were performed on CON data since
290 this was only a subgroup of the total sample and was only used as a
291 reference for the change in liver glycogen concentration with 3 h of exercise in
292 the absence of carbohydrate ingestion. Accordingly, all other comparisons
293 were made between GLU and SUC only.

295 Blood, plasma and respiratory variables and subjective ratings were assessed
296 by two-way (trial x time) repeated measures ANOVA with trial (GLU vs SUC)
297 and time (all time points during exercise) as within-subject factors. Liver and
298 muscle glycogen and intramyocellular lipid concentrations were also assessed
299 by two-way (trial x time) repeated measures ANOVA with trial (GLU vs SUC)

300 and time (pre- vs post-exercise) as within-subject factors. Mean exercise
301 responses in GLU and SUC trials (carbohydrate intake, heart rate, fluid intake
302 and power output) were assessed by paired t-tests. All *P* values are corrected
303 for multiple comparisons (Holm-Sidak). A *P* value of ≤ 0.05 was used to
304 determine statistical significance. All data were analyzed using Prism v5
305 (GraphPad Software, San Diego, CA).

Results

Participants

Participants' characteristics are provided in **Table 2**. No differences were observed for age, body mass, height, $\dot{V}O_{2\text{ peak}}$, W_{peak} , body mass index, systolic or diastolic blood pressure between participants who completed the main trials (GLU and SUC) and the subgroup of participants who also completed the additional CON trial.

Subjective ratings

RPE increased during exercise (time effect, $P < 0.001$), but to less of an extent during SUC when compared to GLU (interaction effect, $P < 0.05$; **Figure 1A**), becoming significantly different between trials from 150 min onwards ($P < 0.05$). Similarly, ratings of gut discomfort increased throughout exercise (time effect, $P < 0.001$) but to less of an extent during SUC when compared to GLU (interaction effect, $P < 0.01$), becoming significantly different at 180 min (Figure 1B, $P < 0.05$).

Respiratory data and whole-body substrate utilization

$\dot{V}O_2$ and $\dot{V}CO_2$ remained stable during exercise (time effect, $P > 0.05$ for both) and were not different between GLU and SUC (both $P > 0.05$). Respiratory exchange ratio (RER) was higher with SUC vs GLU (trial effect, $P < 0.05$) for time points 90 min onwards (interaction effect, $P < 0.05$; **Figure 2C**). Whole-body carbohydrate utilization rates were higher during SUC (2.03 ± 0.43 g/min) when compared with GLU (1.66 ± 0.36 g/min; $P < 0.05$), at the

expense of fat oxidation rates (SUC: 0.35 ± 0.15 vs GLU: 0.48 ± 0.12 g/min; $P < 0.05$), resulting in energy expenditure rates that did not differ between trials (SUC: 8.8 ± 1.2 vs GLU: 8.6 ± 0.9 MJ; $P > 0.05$; **Figure 3A**). In the subgroup who also completed the CON trial ($n = 4$), whole-body fat oxidation rates were lower during both GLU (0.42 ± 0.10 g/min) and SUC (0.33 ± 0.11 g/min), compared to CON (0.64 ± 0.19 g.min; $P < 0.05$), whilst carbohydrate oxidation rates (SUC: 2.04 ± 0.40 vs GLU: 1.79 ± 0.43 vs CON: 1.20 ± 0.44 MJ) did not significantly differ between trials ($P > 0.05$). Accordingly, energy expenditure (SUC: 8.7 ± 0.6 vs GLU: 8.6 ± 0.8 vs CON: 8.4 ± 0.4 MJ) also did not differ between trials ($P > 0.05$; Figure 3B).

Circulating metabolite and insulin concentrations

Blood glucose and plasma insulin concentrations were not significantly different between trials (trial effect, $P > 0.05$; interaction effect, $P > 0.05$ for both variables; **Figure 4A**). In contrast, blood lactate concentrations were higher with SUC vs GLU (trial effect, $P < 0.01$), rising at the onset of exercise (time effect, $P < 0.001$) to a greater extent in SUC vs GLU until 120 min (interaction effect, $P < 0.01$; Figure 4B). Plasma NEFA concentrations fell from ~ 0.5 mmol/L to ~ 0.2 mmol/L during the first hour of exercise before rising again (time effect, $P < 0.001$), the latter of which occurred to a greater degree in GLU compared to SUC (interaction effect, $P < 0.01$; Figure 4D).

Muscle and liver glycogen concentration

Muscle and liver glycogen concentrations are displayed in **Figures 5A, 5B, 5C and 5D**. Pre-exercise, no differences were observed in liver and muscle

glycogen concentrations between trials $P > 0.05$ for both variables). The day-to-day coefficients of variation for pre-exercise liver and muscle glycogen concentrations were 12% and 20%, respectively. The between subject coefficient of variation for pre-exercise liver and muscle glycogen were 54% and 41%, respectively. In the subgroup who also completed the CON trial ($n = 4$), liver glycogen concentrations declined during exercise in CON, but not when either glucose or sucrose were ingested (interaction effect, $P < 0.05$; Figure 5B). In contrast to the liver, muscle glycogen concentrations declined during exercise regardless of trial (trial effect, $P > 0.05$; time effect, $P < 0.01$; interaction effect, $P > 0.05$; Figure 5D).

Post-exercise liver glycogen concentrations did not differ from pre-exercise values when either glucose or sucrose were ingested (time effect, $P > 0.05$; interaction effect, $P > 0.05$). The change in liver glycogen concentrations from pre- to post-exercise was positive with glucose (20 ± 55 mmol/L) and sucrose (27 ± 58 mmol/L; $P > 0.05$ GLU vs SUC) ingestion, but negative in the CON treatment (-171 ± 73 mmol/L).

Muscle glycogen concentrations were reduced following exercise (time effect, $P < 0.001$). The changes in muscle glycogen concentrations did not differ between trials (trial effect, $P > 0.05$; interaction effect, $P > 0.05$; Figures 5C and 4D). The pre- to post-exercise changes in muscle glycogen concentration did not differ between GLU (-40 ± 37 mmol/L) and SUC (-47 ± 36 ; $P > 0.05$).

Intramyocellular lipid concentration

No differences were observed in pre-exercise IMCL concentration ($P > 0.05$) between trials. The day-to-day coefficient of variation for pre-exercise IMCL concentration was 21%. The between-subject coefficient of variation for pre-exercise IMCL concentration was 47%. In the full sample ($n = 14$) exercise decreased IMCL concentrations (time effect $P < 0.01$) to a similar extent in both trials (trial effect, $P > 0.05$; interaction effect, $P > 0.05$; Figure 5E). The pre- to post-exercise changes in IMCL concentration did not differ between GLU ($-1.5 \pm 6.0 \mu\text{mol/g}$) and SUC ($-1.6 \pm 6.4 \mu\text{mol/g}$; $P > 0.05$).

In the subgroup who completed the CON trial ($n = 4$), post-exercise IMCL concentrations were not significantly different to pre-exercise values (time effect, $P > 0.05$), and the responses were not significantly different between trials (trial effect, $P > 0.05$; interaction effect, $P > 0.05$; Figure 5F).

Discussion

In the present study we provide novel data demonstrating that carbohydrate ingestion during endurance type exercise can prevent liver glycogen depletion, and that this effect is independent of the type of carbohydrate (glucose or sucrose) ingested. In contrast, neither glucose nor sucrose ingestion at 1.7 g/min (102 g/h) could attenuate the decline in muscle glycogen following exercise. Sucrose ingestion increased whole-body carbohydrate utilization when compared with glucose ingestion.

Muscle glycogen and plasma glucose are the main fuel sources during prolonged, moderate-intensity endurance type exercise (41). Plasma glucose is maintained during exercise by glycogenolysis and gluconeogenesis, primarily from the liver. Accordingly, continuous exercise lasting more than 60 min substantially depletes liver glycogen concentrations (37). Given that liver glycogen strongly associates with endurance capacity (6), maintaining liver glycogen concentrations is likely to benefit endurance performance. Previous research using glucose tracers has indicated that high rates of glucose ingestion can suppress endogenous glucose appearance (21), implying that carbohydrate ingestion during exercise may attenuate exercise induced liver glycogen depletion. Here we present the first quantitative evidence of liver glycogen maintenance following carbohydrate ingestion during exercise. We found that 3 h of cycling, in the absence of carbohydrate ingestion reduces liver glycogen concentrations by ~49%, which is consistent with previous findings (37). When ingesting ~1.7 g/min (~102 g/h) glucose or sucrose, liver

glycogen concentrations are not lowered during prolonged exercise (Figures 5A and 5B).

Liver glycogen concentrations displayed a relatively high variability between subjects (coefficient of variation: 54%), compared to the day-to-day variability within subjects (coefficient of variation: 12%). This provides an explanation for the relatively higher baseline liver glycogen concentrations in the subgroup that completed the CON trial ($n = 4$; Figure 5B) compared to the entire sample ($n = 14$; Figure 5A).

Carbohydrate ingestion during exercise increases exogenous carbohydrate oxidation and has been shown to spare net muscle glycogen utilization under some conditions (36), although not typically during the latter stages of more prolonged (> 1 h), cycling exercise. These responses are thought to contribute to the performance benefits of carbohydrate ingestion during prolonged exercise (7). The present data demonstrate that neither the ingestion of glucose nor sucrose are able to attenuate net muscle glycogen utilization during prolonged moderate-intensity cycling, even when large quantities of multiple transportable carbohydrate (~ 1.7 g/min; 102 g/h) are ingested that augment exogenous carbohydrate availability. In contrast, whole body carbohydrate utilization rates were higher with sucrose vs glucose ingestion, with a concomitant reduction in fat use. Data from the subgroup also demonstrate that both glucose and sucrose ingestion suppress fat utilization relative to CON, although the numerical differences in carbohydrate utilization rates did not reach statistical significance with the subgroup ($n = 4$; $P = 0.07$).

At rest, fructose is preferentially stored as liver glycogen rather than muscle glycogen. This has led some to speculate that sucrose, when compared with glucose ingestion may be particularly effective at maintaining or increasing liver glycogen during exercise. In the present study, sucrose ingestion did not preserve liver glycogen concentrations to any greater extent than glucose ingestion. In line with previous observations of substantial declines in endogenous glucose production during exercise when glucose was ingested (21), our data seem to suggest that liver glycogen contents are maintained during exercise when ingesting large amounts (~1.7 g/min; ~102 g/h) of glucose or sucrose. The surplus carbohydrates are shunted towards oxidation rather than storage, at the expense of lipid oxidation.

The increase in whole-body carbohydrate utilization following sucrose vs glucose ingestion seems to confirm that sucrose ingestion increases exogenous carbohydrate availability and carbohydrate flux. This shift in metabolism is likely due to a number of coordinated factors, including the higher lactate concentrations observed following sucrose ingestion. Higher circulating lactate concentrations are very likely due to the fructose component of sucrose, the majority of which is converted to lactate and glucose upon bypassing the liver. Glucose-fructose co-ingestion during exercise has been shown to increase plasma lactate and glucose turnover and oxidation (24), with a minimal amount of fructose being directly oxidized (24). The greater whole-body carbohydrate utilization rate following sucrose ingestion is therefore likely attributed to a combination of (greater) plasma lactate, glucose and (to a lesser extent) fructose oxidation rates. Lactate also

inhibits adipocyte lipolysis via the G-protein coupled receptor GPR81 (27). This is likely one of the factors responsible for the lower plasma NEFA concentrations following sucrose versus glucose ingestion in the presence of similar insulinemia. As there were no differences in muscle lipid content changes between treatments, the greater fat use in the glucose compared with the sucrose trial is likely entirely attributed to greater uptake and oxidation of plasma derived NEFA.

Lactate formation is associated with hydrogen ion production, which may displace CO_2 from bicarbonate stores with consequent implications for estimates of $\dot{V}\text{CO}_2$, RER and substrate utilization (13). The ~ 0.5 mmol/L increase in lactate concentration following SUC vs GLU however, would have a negligible (<0.07 mL/min) effect on CO_2 displacement (13). Therefore, values obtained from expiratory gas samples are likely to be a valid representation of net substrate utilization.

We observed a lower RPE towards the end of exercise following sucrose compared with glucose ingestion. This is in spite of the higher lactate concentrations following ingestion of sucrose compared to glucose, offering additional evidence of the disassociation between lactate concentrations and RPE (29). Exogenous carbohydrate oxidation rates have been shown to correlate with exercise performance during prolonged, moderate-to-high intensity exercise (35). Therefore, the lower RPE following sucrose versus glucose ingestion may be attributed to the greater exogenous carbohydrate uptake and subsequent oxidation rates when co-ingesting fructose (17, 19).

493 The lower RPE may of course also be directly attributed to the lesser
494 occurrence of gastrointestinal discomfort when ingesting large amounts of
495 multiple transportable carbohydrates versus glucose only (Figure 1).

496

497 In conclusion, ingestion of large amounts [~ 1.7 g/min (~ 102 g/h), relative to
498 the ~ 1.5 g/min (90 g/h) recommended for exercise lasting >2.5 h] of glucose
499 or sucrose during prolonged endurance type exercise prevent the exercise-
500 induced decline in liver glycogen content without modulating muscle glycogen
501 depletion.

Acknowledgements

The authors wish to thank the volunteers for their time and effort in participating in this study. We also thank R Veasey and J Forster for blinding and test-drink preparation, and Louise Ward, Dorothy Wallace and Tim Hodgson for assistance with magnetic resonance examinations. Professor Trenell is supported by a Senior Fellowship from the National Institute for Health Research.

Disclosures

This study was funded by Sugar Nutrition UK and Suikerstichting Nederland.

512 References

- 513 1. **Betts JA, and Thompson D.** Thinking outside the bag (not necessarily
514 outside the lab). *Med Sci Sports Exerc* 44: 2040; author reply 2041, 2012.
- 515 2. **Borg GA.** Perceived exertion: a note on "history" and methods. *Med*
516 *Sci Sports* 5: 90-93, 1973.
- 517 3. **Bosch AN, Dennis SC, and Noakes TD.** Influence of carbohydrate
518 ingestion on fuel substrate turnover and oxidation during prolonged exercise.
519 *J Appl Physiol* (1985) 76: 2364-2372, 1994.
- 520 4. **Bottomley PA.** Spatial localization in NMR spectroscopy in vivo. *Ann*
521 *New York Acad Sci* 508: 333-348, 1987.
- 522 5. **Burke LM, Hawley JA, Wong SH, and Jeukendrup AE.**
523 Carbohydrates for training and competition. *J Sports Sci* 29 Suppl 1: S17-27,
524 2011.
- 525 6. **Casey A, Mann R, Banister K, Fox J, Morris PG, Macdonald IA, and**
526 **Greenhaff PL.** Effect of carbohydrate ingestion on glycogen resynthesis in
527 human liver and skeletal muscle, measured by (13)C MRS. *Am J Physiol*
528 *Endocrinol Metab* 278: E65-75, 2000.
- 529 7. **Cermak NM, and van Loon LJ.** The use of carbohydrates during
530 exercise as an ergogenic aid. *Sports Med* 43: 1139-1155, 2013.
- 531 8. **Coyle EF, Coggan AR, Hemmert MK, and Ivy JL.** Muscle glycogen
532 utilization during prolonged strenuous exercise when fed carbohydrate. *J Appl*
533 *Physiol* 61: 165-172, 1986.
- 534 9. **Douglas CG.** A method for determining the total respiratory exchange
535 in man. *J Physiol* 42: xvii-xviii, 1911.
- 536 10. **Faul F, Erdfelder E, Lang AG, and Buchner A.** G*Power 3: a flexible
537 statistical power analysis program for the social, behavioral, and biomedical
538 sciences. *Beh Res Methods* 39: 175-191, 2007.
- 539 11. **Ferraris RP, and Diamond J.** Regulation of intestinal sugar transport.
540 *Physiol Rev* 77: 257-302, 1997.
- 541 12. **Flynn MG, Costill DL, Hawley JA, Fink WJ, Neufer PD, Fielding RA,**
542 **and Sleeper MD.** Influence of selected carbohydrate drinks on cycling
543 performance and glycogen use. *Med Sci Sports Exerc* 19: 37-40, 1987.
- 544 13. **Frayn KN.** Calculation of substrate oxidation rates in vivo from
545 gaseous exchange. *J Appl Physiol* 55: 628-634, 1983.
- 546 14. **Gray GM, and Ingelfinger FJ.** Intestinal absorption of sucrose in man:
547 interrelation of hydrolysis and monosaccharide product absorption. *J Clin*
548 *Invest* 45: 388-398, 1966.
- 549 15. **Hargreaves M, and Briggs CA.** Effect of carbohydrate ingestion on
550 exercise metabolism. *J Appl Physiol* 65: 1553-1555, 1988.
- 551 16. **Izumida Y, Yahagi N, Takeuchi Y, Nishi M, Shikama A, Takarada A,**
552 **Masuda Y, Kubota M, Matsuzaka T, Nakagawa Y, Iizuka Y, Itaka K,**
553 **Kataoka K, Shioda S, Niiijima A, Yamada T, Katagiri H, Nagai R, Yamada**
554 **N, Kadowaki T, and Shimano H.** Glycogen shortage during fasting triggers
555 liver-brain-adipose neurocircuitry to facilitate fat utilization. *Nat Commun* 4:
556 2316, 2013.
- 557 17. **Jentjens RL, and Jeukendrup AE.** High rates of exogenous
558 carbohydrate oxidation from a mixture of glucose and fructose ingested during
559 prolonged cycling exercise. *Br J Nutr* 93: 485-492, 2005.

18. **Jentjens RL, Shaw C, Birtles T, Waring RH, Harding LK, and Jeukendrup AE.** Oxidation of combined ingestion of glucose and sucrose during exercise. *Metabolism* 54: 610-618, 2005.
19. **Jeukendrup AE.** Carbohydrate and exercise performance: the role of multiple transportable carbohydrates. *Curr Opin Clin Nutr Metab Care* 13: 452-457, 2010.
20. **Jeukendrup AE, Raben A, Gijsen A, Stegen JH, Brouns F, Saris WH, and Wagenmakers AJ.** Glucose kinetics during prolonged exercise in highly trained human subjects: effect of glucose ingestion. *J Physiol* 515 (Pt 2): 579-589, 1999.
21. **Jeukendrup AE, Wagenmakers AJ, Stegen JH, Gijsen AP, Brouns F, and Saris WH.** Carbohydrate ingestion can completely suppress endogenous glucose production during exercise. *Am J Physiol* 276: E672-683, 1999.
22. **Jeukendrup AE, and Wallis GA.** Measurement of substrate oxidation during exercise by means of gas exchange measurements. *Int J Sports Med* 26 Suppl 1: S28-37, 2005.
23. **Kuipers H, Verstappen FT, Keizer HA, Geurten P, and van Kranenburg G.** Variability of aerobic performance in the laboratory and its physiologic correlates. *Int J Sports Med* 6: 197-201, 1985.
24. **Lecoultre V, Benoit R, Carrel G, Schutz Y, Millet GP, Tappy L, and Schneider P.** Fructose and glucose co-ingestion during prolonged exercise increases lactate and glucose fluxes and oxidation compared with an equimolar intake of glucose. *Am J Clin Nutr* 92: 1071-1079, 2010.
25. **Lim EL, Hollingsworth KG, Smith FE, Thelwall PE, and Taylor R.** Effects of raising muscle glycogen synthesis rate on skeletal muscle ATP turnover rate in type 2 diabetes. *Am J Physiol Endocrinol Metab* 301: E1155-1162, 2011.
26. **Lim EL, Hollingsworth KG, Smith FE, Thelwall PE, and Taylor R.** Inhibition of lipolysis in Type 2 diabetes normalizes glucose disposal without change in muscle glycogen synthesis rates. *Clin Sci* 121: 169-177, 2011.
27. **Liu C, Wu J, Zhu J, Kuei C, Yu J, Shelton J, Sutton SW, Li X, Yun SJ, Mirzadegan T, Mazur C, Kamme F, and Lovenberg TW.** Lactate inhibits lipolysis in fat cells through activation of an orphan G-protein-coupled receptor, GPR81. *J Biol Chem* 284: 2811-2822, 2009.
28. **Macauley M, Smith FE, Thelwall PE, Hollingsworth KG, and Taylor R.** Diurnal variation in skeletal muscle and liver glycogen in humans with normal health and Type 2 diabetes. *Clin Sci* 128: 707-713, 2015.
29. **Miller BF, Fattor JA, Jacobs KA, Horning MA, Navazio F, Lindinger MI, and Brooks GA.** Lactate and glucose interactions during rest and exercise in men: effect of exogenous lactate infusion. *J Physiol* 544: 963-975, 2002.
30. **Mithieux G, Misery P, Magnan C, Pillot B, Gautier-Stein A, Bernard C, Rajas F, and Zitoun C.** Portal sensing of intestinal gluconeogenesis is a mechanistic link in the diminution of food intake induced by diet protein. *Cell Metab* 2: 321-329, 2005.
31. **Naressi A, Couturier C, Castang I, de Beer R, and Graveron-Demilly D.** Java-based graphical user interface for MRUI, a software package for quantitation of in vivo/medical magnetic resonance spectroscopy signals. *Comp Biol Med* 31: 269-286, 2001.

32. **Nilsson LH, and Hultman E.** Liver and muscle glycogen in man after glucose and fructose infusion. *Scand J Clin Lab Invest* 33: 5-10, 1974.
33. **Rossmann A, Butzenlechner M, and Schmidt HL.** Evidence for a nonstatistical carbon isotope distribution in natural glucose. *Plant Physiol* 96: 609-614, 1991.
34. **Rothman DL, Shulman RG, and Shulman GI.** N.m.r. studies of muscle glycogen synthesis in normal and non-insulin-dependent diabetic subjects. *Biochem Soc Trans* 19: 992-994, 1991.
35. **Smith JW, Zachwieja JJ, Peronnet F, Passe DH, Massicotte D, Lavoie C, and Pascoe DD.** Fuel selection and cycling endurance performance with ingestion of [13C]glucose: evidence for a carbohydrate dose response. *J Appl Physiol* 108: 1520-1529, 2010.
36. **Stellingwerff T, Boon H, Gijzen AP, Stegen JH, Kuipers H, and van Loon LJ.** Carbohydrate supplementation during prolonged cycling exercise spares muscle glycogen but does not affect intramyocellular lipid use. *Pflugers Archiv : Eur J Physiol* 454: 635-647, 2007.
37. **Stevenson EJ, Thelwall PE, Thomas K, Smith F, Brand-Miller J, and Trenell MI.** Dietary glycemic index influences lipid oxidation but not muscle or liver glycogen oxidation during exercise. *Am J Physiol Endocrinol Metab* 296: E1140-1147, 2009.
38. **Tsintzas K, and Williams C.** Human muscle glycogen metabolism during exercise. Effect of carbohydrate supplementation. *Sports Med* 25: 7-23, 1998.
39. **Tsintzas K, Williams C, Constantin-Teodosiu D, Hultman E, Boobis L, Clarys P, and Greenhaff P.** Phosphocreatine degradation in type I and type II muscle fibres during submaximal exercise in man: effect of carbohydrate ingestion. *J Physiol* 537: 305-311, 2001.
40. **Tsintzas OK, Williams C, Boobis L, and Greenhaff P.** Carbohydrate ingestion and glycogen utilization in different muscle fibre types in man. *J Physiol* 489 (Pt 1): 243-250, 1995.
41. **van Loon LJ, Greenhaff PL, Constantin-Teodosiu D, Saris WH, and Wagenmakers AJ.** The effects of increasing exercise intensity on muscle fuel utilisation in humans. *J Physiol* 536: 295-304, 2001.
42. **Vandenbogaerde TJ, and Hopkins WG.** Effects of acute carbohydrate supplementation on endurance performance: a meta-analysis. *Sports Med* 41: 773-792, 2011.
43. **Wallis GA, and Wittekind A.** Is there a specific role for sucrose in sports and exercise performance? *Int J Sport Nutr Exerc Metab* 23: 571-583, 2013.

FIGURE LEGENDS

Figure 1 Ratings of perceived exertion (A) and gut discomfort (B) during 3 h of cycling with ingestion of glucose or sucrose in trained cyclists ($n = 14$). Data are expressed as means \pm 95% CI. * $P < 0.05$, significantly different between GLU and SUC. GLU, glucose; SUC, sucrose.




Figure 2 VO_2 (A), VCO_2 (B) and respiratory exchange ratio (C) during 3 h of cycling with ingestion of glucose or sucrose in trained cyclists ($n = 14$). Data are expressed as means \pm 95% CI. * $P < 0.05$, significantly different between GLU and SUC. GLU, glucose; SUC, sucrose; VCO_2 , rate of carbon dioxide production; VO_2 , rate of oxygen consumption.




Figure 3 Substrate utilization during 3 h of cycling with ingestion of glucose or sucrose in trained cyclists (A; $n = 14$) and in the subgroup of trained cyclists (B; $n = 4$). Data are expressed as means \pm 95% CI. * $P < 0.05$, significantly different from CON. CHO, carbohydrate; GLU, glucose; SUC, sucrose; CON, water control.




Figure 4 Blood glucose (A) and lactate (B), and plasma insulin (C) and NEFA (D) concentrations during 3 h of cycling with ingestion of glucose or sucrose in trained cyclists ($n = 14$). Data are expressed as means \pm 95% CI. * $P < 0.05$, significantly different between GLU and SUC. GLU, glucose; NEFA, non-esterified fatty acid; SUC, sucrose.




Figure 5 Liver glycogen (A), muscle glycogen (C) and intramyocellular lipid (E) concentrations prior to (Pre) and immediately following (Post) 3 h of cycling with ingestion of glucose or sucrose in the full sample of trained cyclists ($n = 14$), and in the subgroup of trained cyclists (B, D and F; $n = 4$). Data are expressed as means \pm 95% CI. * $P < 0.05$, significantly different when compared with pre-exercise values. GLU, glucose; SUC, sucrose; CON, water control.




Table 1 Carbohydrate intake and physiological variables of trained cyclists during 3 h of cycling with ingestion of glucose, sucrose or water.

| | GLU (<i>n</i> = 14) | SUC (<i>n</i> = 14) | CON (<i>n</i> = 4) |
|-----------------------------|-------------------------|-------------------------|------------------------|
| Carbohydrate intake (g/min) | 1.7 ± 0.2 | 1.7 ± 0.2 | 0 ± 0 |
| Fluid intake (L) | 2.1 ± 0.2 | 2.2 ± 0.2 | 2.3 ± 0.0 |
| Power output (W) | 165 ± 17 | 165 ± 17 | 158 ± 7 |
| Mean heart rate (beats/min) | 145 ± 14 | 146 ± 12 | 122 ± 8 |

Data are expressed as means ± SEM. GLU, glucose; SUC, sucrose; CON, water control.

Table 2 Characteristics of trained cyclists who completed GLU, SUC and CON trials.

| | GLU and SUC | CON |
|---------------------------------------|------------------|-----------------|
| | (<i>n</i> = 14) | (<i>n</i> = 4) |
| Age (y) | 25 ± 5 | 26 ± 6 |
| Body mass (kg) | 73.1 ± 9.3 | 75.3 ± 10.7 |
| Height (m) | 1.78 ± 0.08 | 1.75 ± 0.09 |
| $\dot{V}O_{2\text{peak}}$ (mL/min/kg) | 58 ± 5 | 60 ± 7 |
| W_{peak} (W) | 330 ± 35 | 316 ± 27 |
| BMI (kg/m ²) | 23.0 ± 1.9 | 24.5 ± 1.8 |
| Systolic blood pressure | 133 ± 11 | 129 ± 6 |
| (mmHg) | | |
| Diastolic blood pressure | 74 ± 8 | 71 ± 8 |
| (mmHg) | | |

Data are expressed as means ± SD. GLU, glucose; SUC, sucrose; $\dot{V}O_{2\text{peak}}$, peak oxygen uptake; CON, water control; W_{peak} , peak power output.